

October 30, 2019

To whom it may concern:

Company Name Corporate Representative	TOHO HOLDINGS CO., LTD. Norio Hamada, Chairman of the Board and Chief Executive Officer (CEO), Representative Director (First Section of Tokyo Stock Exchange Securities Code:8129)
Contact:	Makoto Kawamura, Director and General Manager, Public and Investor Relations Department (TEL: 81-3-6838-2803)

**Notice Regarding Approval of Partial Amendments to Dosage and Administration of
Pitavastatin Calcium Tablets 1mg / 2mg “FFP” for KYOSOMIRAI PHARMA**

TOHO HOLDINGS CO., LTD. (Headquarters: Tokyo; Chairman of the Board and Chief Executive Officer (CEO), Representative Director: Norio Hamada) is pleased to announce that KYOSOMIRAI PHARMA CO., LTD. (Headquarters: Tokyo; President and Representative Director: Atsushi Udoh), the wholly-owned subsidiary involved in pharmaceuticals manufacture and sales business, received an approval of partial amendments to dosage and administration of Pitavastatin Calcium Tablets 1mg / 2mg “FFP” on October 30, 2019 as below.

Outline of Pitavastatin Calcium Tablets 1mg / 2mg “FFP”

Class	Product Name	Original Brand Name
HMG-CoA Reductase Inhibitor	Pitavastatin Calcium Tablets 1mg / 2mg “FFP”	LIVALO Tablets 1mg / 2mg

Outline of Amendments to Dosage and Administration of Pitavastatin Calcium Tablets 1mg / 2mg “FFP”

Indications and Usage	Dosage and Administration
Hypercholesterolemia	<p>The usual oral dose for adults is 1-2mg of pitavastatin calcium once daily. The dosage may be adjusted according to the patient’s age and symptoms. If lowering of LDL cholesterol level is insufficient, the dosage can be increased up to 4mg daily.</p>
Familial hypercholesterolemia	<p>Adults: The usual oral dose for adults is 1-2mg of pitavastatin calcium once daily. The dosage may be adjusted according to the patient’s age and symptoms. If lowering of LDL cholesterol level is insufficient, the dosage can be increased up to 4mg daily.</p> <p><u>Children:</u> <u>The usual oral dose for children aged 10 years and older is 1mg of pitavastatin calcium once daily.</u> <u>The dosage may be adjusted according to the patient’s symptoms. If lowering of LDL cholesterol level is insufficient, the dosage can be increased up to 2mg daily.</u></p>

(Underlined parts are revised parts.)

< Reference >

Adults

	Tablets 1mg	Tablets 2mg	Tablets 4mg
Hypercholesterolemia	○	○	○
Familial hypercholesterolemia	○	○	○

Children

	Tablets 1mg	Tablets 2mg	Tablets 4mg
Hypercholesterolemia	—	—	—
Familial hypercholesterolemia	○	○	—

○ : Approved dosage and administration — : Not approved